4267 FIRST AID KIT- 4267 first aid kit Honeywell Safety Products USA, INC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

4267 First Aid Kit (Neomycin, EW, ammonia inh, alcohol wipes, Burn Sray, Antiseptic Spray, Hand Sanitizer, aypanal- FAK3SHLF-CLSB)

Eyes aline

Active ingredient

Sterile Water 99%

Eyes aline

Purpose

Eyewash

Eyes aline

Uses

• for flushing the eye to remove loose foreign material, air pollutants or chlorinated water

Eyes aline

Warnings

For external use only-

Obtain immediate medical treatment for all open wounds in or near eyes.

To avoid contamination, do not touch tip of container to any surface.

Do not reuse. Once opened, discard.

Do not use

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away.

Stop use and ask a doctor if

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Eyes aline

Directions

remove contacts before using

- twist top to remove
- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

Eyes aline

Inactive ingredients

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

Eyes aline *Questions*

1-800-430-5490 Honeywell Sadety Products USA, Inc. Smithfield, RI 02917

Alcohol Wipe Active ingredient

Isopropyl alcohol 70%

Alcohol Wipe Purpose

First aid antiseptic

Alcohol Wipe

Uses

• first aid to help prevent infection in minor cuts, scrapes, and burns

Alcohol Wipe Warnings

For external use only

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burn

When using this product

do not use longer than one week unless directed by a doctor

Stop use and consult a doctor

• if condition persists or gets worse

Keep out of reach of children

• If swallowed, get medical help or contact a Poison Control Center right away.

Alcohol Wipe

Directions

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

Alcohol Wipe Other information

store at room temperature 15 0 to 25 0 C (59 0 to 77 0 F)

Alcohol Wipe Inactive ingredient

water

Alcohol Wipe Questions

1-800-430-5490

Aypanal

Active ingredient

Acetaminophen 325 mg

Aypanal Purpose

Pain reliever/ fever reducer

Aypanaly

Uses

• temporarily relieves minor aches and pains due to the common cold and headache - temporarily reduces fever

Keep out of reach of children.

Keep out of reach of children.

Aypanal

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen

• adult has 3 or more alcoholic drinks every day while using this product:

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin rash occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if

• you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

Overdose warning

- In case of accidental overdose, get medical help or contact a Poison Control Center right away.
- Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Aypanal

Directions

do not take more than directed (see overdose warning)

ladults and children 12 years of age or older

- take two tablets every 4-6 hours while symptoms last
- do not take more than 12 tablets in 24 hours

children 6 to under 12 years of age

- take 1 tablet every 4-6 hours while symptoms last
- do not take more than 5 tablets in 24 hours

children under 6 years consult a doctor

Aypanal

Other information

- store at room temperature 15 0 to 30 0 C (59 0 86 0 F)
- TAMPER EVIDENT PACKETS- DO NOT USE IF OPEN OR TORN

Aypanal

Inactive ingredients

corn starch, microcrystalline cellulose, povidone, sodium starch glycolate, stearic acid

Aypanal Questions

1-800-430-5490

Neomycin Active ingredient

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

Neomycin

Purpose

First aid antibiotic

Neomycin

Uses

• first aid to help prevent infection in - minor cuts - scrapes - burns

Do not use

- in the eyes
- over large areas of the body

Neomycin

Warnings

For external use only

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- a rash or other allergic reaction develops
- you need to use longer than 1 week

Stop use and ask a doctor if

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of reach of children

• If swallowed, get medical help or contact a Poison Control Center right away

Neomycin

Direction

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Neomycin Other information

store at 15 0 to 25 0 C (59 0 to 77 0 F)

Neomycin Inactive ingredient

petrolatum

Neomycin *Questions?*

1-800-430-5490

Antis eptic Spray *Active ingredient*

Benzalkonium chloride 0.13%

Antis eptic Spray *Purpose*

First aid antiseptic

Antis eptic Spray

Uses

• first aid to help prevent infection in minor cuts, scrapes and burns

Antiseptic Spray

Warnings

For external use only

Do not use

- in or near the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

When using this product

• do not use longer than one week unless directed by a doctor

Stop use and ask a doctor if

• the condition persists or gets worse

Keep out of reach of children

• If swallowed, get medical help or contact a Poison Control Center right away

Antiseptic Spray

Directions

- clean the affected area
- spray a small amount of this product on the area 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

Antiseptic Spray

Other information

- shake well
- store at room temperature $15^{0}-30^{0}$ C ($59^{0}-86^{0}$ F)

Antiseptic Spray Inactive ingredients

diazolidinyl urea, edetate disodium, glycerin, hypromellose, methylparaben, octoxynol 9, propylene glycol, propylparaben, trolamine, water

Antiseptic Spray Questions

1-800-430-5490

Burn Spray Active ingredient

Benzethonium chloride 0.2% w/w

Benzocaine 10% w/w

Menthol 0.33% w/w

Burn Spray

Purpose

Benzethonium chloride 0.2% w/w

Benzocaine 10% w/w

Menthol 0.33% w/w

Burn Spray

<u>Uses</u>

for the temporary relief of pain and itching and helps protect against infection in:

- minor cuts and scrapes
- burns
- sunburn
- insect bites
- minor skin irritations

Burn Spray Warnings

For external use only

Flammable

- keep away from fire or flame
- contents under pressure
- do not puncture or incinerate container
- do not expose to temperatures above 120 0 F

Do not use

- in or near the eyes or other mucous membranes
- in case of serious burns
- in case of deep or puncture wounds
- for prolonged period of time
- on large portion of the body

Stop use and ask a doctor if

- condition worsens or symptoms persist for more than 7 days
- condition clears up and recurs within a few days
- redness, swelling, or irritation occurs

Keep out of the reach of children

• If swallowed, get medical help or contact a Poison Control Center right away.

Burn Spray

Directions

- clean the affected area
- shake can well before using
- hold 4 6 inches from surface and spray area until wet
- may be covered with a sterile bandage, if bandaged let dry first
- for adult institutional use only
- not intended for use on children

Burn Spray

Other information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating or inhaling the contents may be harmful or fatal

Burn Spray

Inactive ingredients

dipropylene glycol, isobutane, n-butane, propane

Hand Sanitizer *Active ingredient*

Ethyl alcohol 62%

Hand Santizer

Purpose

Antiseptic handwash

Hand Sanitizer

Uses

- for hand washing to decrease bacteria on skin
- recommended for repeated use

Hand Sanitizer

Warnings

For external use only

Flammable, keep away from fire or flame

When using this product

- do not use in the eyes
- discontinue use if irritation and redness develops. If condition persists for more than 72 hours consult a doctor.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Hand Sanitizer

Directions

wet hands thoroughly with product and allow to dry without wiping

Hand Sanitizer Other information

store at 15 0 to 25 0 C (59 0 to 77 0 F)

Hand Sanitizer Inactive ingredients

acrylates/C10-30 alkyl acrylate crosspolymer, aloe barbadensis leaf juice, dl-alpha tocopheryl acetate, fragrance, PEG-60 almond glycerides, propylene glycol, purified water, triisopropanolamine

Hand Sanitizer

Questions or Comments?

1-800-275-3433 info@waterjel.com www.waterjel.com

Ammonia

Active ingredient

Ammonia 15%

Ammonia

Purpose

Respiratory stimulant

Ammonia

Uses

• to prevent or treat fainting

Ammonia

Warnings

For external use only

Do not use

• if you have breathing problems such as asthma or emphysema

Stop use and ask a doctor if

• condition persists

Keep out of reach of children

If swallowed get medical help or contact a Poison Control Center right away.

Ammonia

Directions

- hold inhalant away from face and crush ampoule between thumb and forefinger at position indicated on sleeve.
- hold near nostrils for inhalation of volatile vapor

Ammonia

Other information

• store at room temperature away from light

Ammonia

Inactive ingredients

alcohol USP, FD&C red #40, lavender oil, lemon oil fcc, nutmeg oil, purified water

Ammonia Questions or Comments?

1-800-430-5490

Eyes aline Principal Display Panel



Datos de medicamento (Para EE.UU. solamente)
Ingrediente Activo Propósito
Agua esteril 99% Lavaojos

Usos
para el lavado de ojo para quitar las particulas sueltas y extrañas, los contaminantes aeros, o agua de cloruro

Advertencias
Para el uso externo sólo - Obtenga tratamiento médico inmediato para todas las heridas abiertas en o cerca de los ojos. Para evitar la contaminación, no toque la punta del envase con ninguna superficie. No vuelva a usar. Vez abierto, descarte.

No se use - si la solución se enturbia o cambia de color e intenenta dora de color e intenenta dora de color e intenenta dora de jo e cambio de visión e rojez continuo o intración del gio e cambio de visión e rojez continuo o intración del gio e la condición emperor o persiste

Manténgase fuera del alcance de los niños.
En caso de ingestión accidental, obtenga atención médica o llame a un Centro de control de envenenamiento immediatamente.

Instrucciones

quitese los lentes de contacto antes de usar la solución

e uerza la tapa para quitar

e njuague el área afectada según se necesite

o controle el chorro haciendo presión el la botella


s ies necesario, sigue enjuagado con un lavaojos o ducha de emergencia

Ingredientes inacctivos
cloruro de sodio, fosfato de sodio monobásico

¿Preguntas? Llame al 1-800-430-5490

Honeywell Safety Products USA, Inc. Smithfield, Rl. 02917


Information Usages Pour le rinçage des yeux afin d'enlever un corps étranger, des polluants atmospheriques où de l'eau chlorée. Advertissements Pour usage externe seulement - Obtenir immédiatement des soins médicaux pour toutes les plaies ouvertes dans ou près des yeux. Pour éviter toute contamination, ne pas toucher la pointe du récipient à n'importe quelle surface. Ne pas réutiliser. Une fois ouvert, jetez-les. Ne pas utiliser • si la solution a changé de couleur ou si elle est brouillée • si vous avez des plaies ouvertes aux yeux ou à proximité, consultez immédiatement un médecin • vous ressentez une douleur oculaire • si voits evision change • rougeur ou irritation persistainte des yeux • condition empre ou persista. Carder hors de la portée des enfants. En cas d'ingestion, communiquer immédiatement avec un médecin ou avec un centre antipoison. Mode d'emploi • enlever les verres de contact avant l'utilisation • dévisser le besoins • ajuster le débit d'écoulement de la solution en augmentant ou en réduisant la pression exercée sur le contennant s' si nécessaire, continuer de rincer avec unesolution de rinçage coulaire d'ugence ou une douche Ingrédients ea stérile, chlorure de sodium, phosphate dibasique de sodium phosphate monobasique de sodium Des questions? Faites le 1-800-430-5490 Honeywell Safety Products USA, Inc. Smithfield, RI. 02917



Aypanal Principal Display Panel



796041-25 Rev A Unit Carton Printing Plate for "C" size carton.





Burn Spray Principal Display Panel



Cat. No. 201005

DRUG FACTS Active ingredients Purpose Benzethonium chloride 0.2% w/w. Benzocaine 10% w/w. Topical antiseptic Topical anesthetic Topical anesthetic Uses • for the temporary relief of pain and itching and helps to protect against infection in · minor cuts and scrapes · burns · sunburn · insect bites · minor skin irritations Warnings For external use only Flammable • keep away from fire or flame • do not puncture or incinerate container contents under pressure do not expose to temperatures above 120°F Do not use • in or near eyes or other mucus membranes • in case of serious burns in case of deep or puncture wounds . for a prolonged period of time on large portion of the body Stop use and ask a doctor if: · conditions worsens or symptoms persist for more than 7 days · condition clears up and recurs within a few days · redness, swelling or irritation occurs Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Directions • clean the affected area • shake can well before using hold 4-6 inches from surface and spray area until wet may be covered with a sterile bandage. If bandaged, let dry first for adult institutional use only not intended for use on children · use only as directed Other information · avoid inhaling · intentional misuse by deliberately concentrating and inhaling the contents may be harmful Inactive ingredients dipropylene glycol, isobutane, n-butane, propane



Questions or comments? 1-800-430-5490

Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917






Hand Sanitizer

Antiseptic Gel
With Vitamin E & Aloe

Kills 99.9% of Germs Without Water

240mL - (8 fl oz)

Ammonia Principal Display Panel



777028B Rev. A white printed four color process blue (pms 072C) and red (pms 186C)



4267 Kit Contnets FAK3SHLF-CLSB

- 1 1X3 PLASTIC 100/BOX
- 1 FINGERTIP "T" WOVEN 40/BOX
- 1 SWIFT KNUCKLE 40/BX
- 11X3WOVEN 100/BOX
- 3 NEOMYCIN ANTIBIOTIC 10 PER
- 1 AMMONIA INHALANTS 10 PER
- 2 EYE DRESS PKT W/4 ADH STRIPS
- 1 TOURNIQUET, 1 PER
- 1 WIRE SPLINT 1 PER
- 1 ADH BAND, EXTRA LARGE, 6 PER
- 1 ALCOHOL PREP PADS 10P
- 1 O/H PUMP ANTISEPTIC 2 OZ ID F
- 1 O/H PUMP BURN RELIEF 2 OZ ID G
- 1 FIRST AID GUIDE ASHI
- 2 TAPE ADHESIVE 1"X 5 YD PLSTC
- 10 HAND SANITIZER 0.9G WJ BULK

- 4 GAUZE CLEAN-WRAP BDGE N/S 2"
- 4 GAUZE CLEAN-WRAP BDGE N/S 4"
- **4 BLOODSTOPPER**
- 1 NON-ADHERENT PADS 2"X3" 10'S
- 1 GZE PADS STERILE 2"X 2" 10'S
- 1 GZE PADS STERILE 3"X 3" 25'S
- 1 ELASTIC BANDAGE 3" X 4.5YD
- 1 CPR FILTERSHIELD 77-100
- 1 COTTON TIPS 100 PER VIAL
- 1 AYPANAL NON-ASP IND 2/ENV 100
- 1 4OZ BFS EYEWASH TRILINGUAL BOTTLE
- 1 SCISSOR BDGE 4" RED PLS HDL
- 1 KIT TWEEZER 3 1/2" SLANTED
- 1 180 EMPTY BLANK NO LOGO
- 1 POCKET INSERT RED #180 KIT 4R
- 1 TONGUE BLADES SR WRAPPED 6'S
- 1 LBL STOCK 6-3/8"X4"
- 1 LBL STOCK 4"X2-7/8"
- 1 LBL STOCK 3"x1-7/8"
- 4 PR LRG NITRILE GLVES ZIP BAG
- 2 WATER-JEL BURN DRESSING 4 X 4
- 1 LBL CONTENTS ANSI 2015 CL B
- 1 LBL CAB CVR ANSI 2015 CL B
- 2 TRI BNDG NON WOVEN 40"X40"X56"
- 2 COLD PACK UNIT 4"X6" BULK
- 1 RED BIO BAGS 2/BX

4267 FIRST AID KIT

4267 first aid kit kit

_					-		r			. •	
v	MO		\mathbf{m}	rt.	-	m	hn	MI	ทว	tio	n
	ΙU	u	u			ш	w		на	uv	

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0498-4267

P	ac	ka	gi	ng

l	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1	NDC:0498-4267-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

Quan	Quantity of Parts				
Part #	Package Quantity	Total Product Quantity			
Part 1	1 BOTTLE	118 mL			
Part 2	10 POUCH	4 mL			
Part 3	50 PACKET	100			
Part 4	1 BOTTLE, SPRAY	59 mL			
Part 5	1 BOTTLE, SPRAY	59 mL			
Part 6	30 PACKET	27 g			
Part 7	10 PACKAGE	9 mL			
Part 8	10 AMPULE	3 mL			

Part 1 of 8

EYESALINE EMERGENCY EYEWASH

purified water liquid

Product Information	
Item Code (Source)	NDC:0498-0100
Route of Administration	OPHTHALMIC

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	98.6 mL in 100 mL		

Inactive Ingredients				
Ingredient Name	Strength			
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
SODIUM PHO SPHATE, DIBASIC (UNII: GR686LBA74)				
SO DIUM PHO SPHATE, MO NO BASIC, MO NO HYDRATE (UNII: 593YOG76RN)				

l	Packaging						
l	#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
l	1	NDC:0498-0100-02	118 mL in 1 BOTTLE; Type 0: Not a Combination Product				

Marketing Info	Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date			
OTC monograph final	part349	12/18/20 18				

Part 2 of 8

ALCOHOL WIPE

isopropyl alcohol swab

Product Information

Item Code (Source) NDC:0498-0143

Route of Administration TOPICAL

Active Ingredient/Active Moiety

receive resident control of the cont		
Ingredient Name	Basis of Strength	Strength
	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name Strength

WATER (UNII: 059QF0KO0R)

Packaging

l	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

9			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	09/18/2018	

Part 3 of 8

AYPANAL NON-ASPIRIN

acetaminophen tablet

Product Information

Item Code (Source) NDC:0498-2001

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg

Inactive Ingredients		
Ingredient Name	Strength	
STARCH, CORN (UNII: O8232NY3SJ)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
PO VIDO NE (UNII: FZ989 GH94E)		
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)		

Product Characteristics			
Color	white	Score	2 pieces
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	circle;U
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/10/2012	

Part 4 of 8

BURN RELIEF

lidocaine hydrochloride spray

Product Information	
Item Code (Source)	NDC:0498-0221
Route of Administration	TOPICAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LIDO CAINE HYDRO CHLO RIDE (UNII: V13007Z41A) (LIDO CAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	24.64 mg in 1 mL

Inactive Ingredients	
Ingredient Name	Strength
EDETATE DISO DIUM (UNII: 7FLD91C86K)	
GLYCERIN (UNII: PDC6A3C0OX)	

TROLAMINE (UNII: 9O3K93S3TK)	
PROPYLPARABEN (UNII: Z8 IX2SC1OH)	
PROPYLENE GLYCOL (UNII: 6 DC9 Q 16 7 V3)	
WATER (UNII: 059QF0KO0R)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TEA TREE OIL (UNII: VIF565UC2G)	
METHYLPARABEN (UNII: A2I8 C7HI9 T)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	

ı	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:0498-0221- 59	59 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	09/18/2018	

Part 5 of 8

ANTISEPTIC

benzalkonium chloride spray

Product Information	
Item Code (Source)	NDC:0498-0402
Route of Administration	TOPICAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 mL	

Inactive Ingredients	
Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
OCTOXYNOL 9 (UNII: 7JPC6 Y25QS)	
GLYCERIN (UNII: PDC6A3C0OX)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	

EDETATE DISO DIUM (UNII: 7FLD9 1C86K)	
TROLAMINE (UNII: 9O3K93S3TK)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29 V3WO)	

ı	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:0498-0402- 59	59 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	09/18/2018	

Part 6 of 8

NEOMYCIN

antibiotic ointment

Product Information	
Item Code (Source)	NDC:0498-0730
Route of Administration	TOPICAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:116QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g		

Inactive Ingredients		
1	Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)		

l	Pa	ckaging			
l	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
l	1 N	NDC:0498-0730-01	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	03/31/2010	

Part 7 of 8

INSTANT HAND SANITIZER

alcohol liquid

Product Information

Item Code (Source) NDC:59898-420

Route of Administration TOPICAL

TRIISO PRO PANO LAMINE (UNII: W9 EN9 DLM98)

Active Ingredient/Active Moiety

Ingredient Name
Basis of Strength
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)
ALCOHOL (UNII: 3K9958V90M) ALCOHOL 62 mL in 100 mL

Inactive Ingredients

inactive ingredients		
Ingredient Name	Strength	
ALOE VERA LEAF (UNII: ZY8 1Z8 3H0 X)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
.ALPHATO COPHERO L ACETATE, DL- (UNII: WR1WPI7EW8)		
CARBOMER COPOLYMER TYPE A (UNII: 71DD5V995L)		
WATER (UNII: 059QF0KO0R)		

Packaging

1 NDC:59898-420-36 0.9 mL in 1 PACKAGE; Type 0: Not a Combination Product

Marketing Information

Marketing Category Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph not final	part333E	04/15/2010	

Part 8 of 8

AMMONIA INHALENT

ammonia inhalent inhalant

Product Information

Item Code (Source) NDC:0498-3334

Route of Administration RESPIRATORY (INHALATION)

Active Ingredient/Active Moiety

	Ingredient Name	Basis of Strength	Strength	
ı	AMMONIA (LINII: 5138 O 19 F1Y) (AMMONIA _ LINII: 5138 O 19 F1Y)	A MMO NIA	0.045 g in 0.3 mI	

AMMO NIA (UNII: 5138 Q 19 F1X) (AMMO NIA - UNII:5138 Q 19 F1X)

MMO NIA 0.045 g in 0.3 mL

Inactive Ingredients

Ingredient Name	Strength

ALCOHOL (UNII: 3K9958V90M)

Packaging

l	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		NID C 0 400 DDD 4 00			

1 NDC:0498-3334-00 0.3 mL in 1 AMPULE; Type 0: Not a Combination Product

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

Labeler - Honeywell Safety Products USA, INC (079287321)

Establishment

Name	me Address ID/		Business Operations
James Alexander		040756421	manufacture(0498-3334)

Establishment

Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, INC		079287321	pack(0498-4267)

Establishment

Name	Address	ID/FEI	Business Operations
Ultra Seal Corporation		085752004	manufacture(0498-2001)

Establishment

Name	Address	ID/FEI	Business Operations
Water-Jel Technologies		155522589	manufacture(0498-0730, 0498-0402, 0498-0221, 59898-420)

Establishment					
Name	Address	ID/FEI	Business Operations		
Honeywell Safety Products USA, Inc.		167518617	manufacture(0498-0100)		

Establishment						
Name	Address	ID/FEI	Business Operations			
Changzhou Maokang Medical		421317073	manufacture(0498-0143)			

Revised: 4/2019 Honeywell Safety Products USA, INC